

2445. GAS ANALYSER FOR MIXTURES WITH TWO THREE AND FOUR COMPONENTS.
 Kravkov, Yu.B., Kamsolkin, VV and Bashkirov, AN (Izvest.
 Akad. Sci.), Nov. 1949, (11), 1649). A new type of gas
 analyser is recommended for CO₂, CO, H₂ and N₂ (or CH₄).
 The analysis is carried out automatically with gas passing
 through the apparatus at different velocities. If the
 velocity is 12-20 lb/hr the analysis takes only a few
 minutes, but this period increases as the velocity of the
 gas current diminishes. The theoretical principles of the
 method and its accuracy are discussed. The error for a
 binary system should be $\pm 1\%$ and for 3- and 4-component mixtures
 $\pm 1.5\%$. It is explained where and how to apply the gas
 analyser.

BCeramRA

Petroleum Inst, AS USSR

F																										T																									
KAMZOLKIN																										VASHKIROV																									
F																										T																									
4711. APPARATUS FOR MEASURING RATE OF GAS FLOW UNDER PRESSURE																																																			
Zil'berg, G.A., Kamsolkin, V.V., Konstantinov, A.A. and Vashkirov, A.N.																																																			
(Zavod. Lab. (Factory Lab.), 1950, vol. 16, 320).																																																			
458-31A METALLURGICAL LITERATURE CLASSIFICATION																																																			
E-2																																																			

Chem Hydrogenation of hexene over a synthetic-ammonia,
iron-silica catalyst. A. A. Kuchin, B. Z. Kamich,
USSR. *Trudy Khim. Prom. (1964) 42, 147-150.* The capacity of the catalyst to reduce hex-
ene depends on its phase compn. and is detd. by the pres-
ence of metallic Fe in it. Upon oxidation, the catalyst ac-
tivity decs; repeated reduction restores the activity to its
original value. An increase in temp. from 300 to 350°
raises the conversion from 5.2 to 12.9%; a further temp.
inc does not affect the degree of hydrogenation. The use
of H₂ only or of a mixt. of H₂ and CO with the partial pres-
sure of H₂ being equal results in practically the same degree
of conversion. An increase in partial pressure of H₂ from
120 to 180 atm. increases the yield of hexane from 20.3 to
27.7%. A contact time of 5-6 min. results in a yield of
almost 60%; decreasing the contact time to half reduces
the yield to 26%; a further decrease of contact time to 47
sec. reduces the yield to 4.2%. By-products consist of
small amts. of acids, alcoh., and esters. B. Z. Kamich

3

7

PM 24

KAMZOLKIN, V. V.

Chem. Ind.
 Nature of changes of iron catalysts in the synthesis of hydrocarbons. A. N. Baskin, V. V. Kamzolkin, and M. I. Kholodkaya. *Trudy Inst. Khim. Neftekhim. Nats. S.S.S.R.* 6, 147-50 (1965).—Studies were conducted with 2 types of catalysis: (1) an alloy catalyst used in the synthesis of NH_3 , and (2) an Fe catalyst pptd. on kieselguhr. In case of the latter, after a definite period changes in phase compn. of catalyst under the influence of oxidation-reduction agents (CO , H_2 , CO_2 , and H_2O) cease and from then on the phase compn. remains practically unchanged. This was not observed for the alloy catalyst. B. Z. K.

3

DM

Catalytic activity of some natural silicate minerals in the synthesis from carbon monoxide and hydrogen. A. N. Buzik, P. I. Novak, S. M. Loktev, and V. V. Kuznetsov.

Khim. i Tekhnol. Toplen 1956, No. 3, 38-42.

The following minerals were included in the study of their catalytic activity in reaction of CO and H₂ (1:1 mole ratio) at 300-350° at 30 atm. pressure: dumortierite (H₂B₁₀Si₂O₂₂), albite (NaAlSi₃O₈), topaz (Al₂F₂(OH)₂(SiO₃)₂), danburite (CaB₂(SiO₃)₂), rhodonite (Mn,Ca(SiO₃)₂), beryl (Be₃Al₂(SiO₃)₆), diopside (CaMg(SiO₃)₂), hedenbergite (CaFe(SiO₃)₂), aegirite (NaFe(SiO₃)₂), spodumene ((Li,Na)Al(SiO₃)₂), actinolite (Ca₂(Mg,Fe)₅(Si₈O₂₂)(OH)₂), gedrite (?), anthophyllite ((Mg,Fe)₂(SiO₃)₂(OH)₂), kunoite (Al₂(SiO₃)₂(OH)₂), montmorillonite ((Mg,Ca)O·Al₂O₃·5SiO₂·7H₂O), talc (Mg₃(Si₄O₁₀)(OH)₂), phlogopite (KMg₃(Si₄AlO₁₀)(F,OH)₂), lepidolite (K₂Si₄Al₂(AlSi₄O₁₀)(F,OH)₂), biotite (K(Mg,Fe)₂(Si₄AlO₁₀)(F,OH)₂), serpentine (Mg₃(Si₄O₁₀)(OH)₂), nontronite (H₂FeSi₄O₁₀·H₂O), chlorite (H₂(Mg,Fe)₂(Al₂Si₄O₁₀), pyrophyllite (Al₂(Si₄O₁₀)(OH)₂), albite (NaAlSi₃O₈), microcline (KAlSi₃O₈), scapolite oligoclase (NaAlSi₃O₈), pollucite (CsAlSi₃O₈), petalite ((Li,Na)AlSi₃O₈), nepheline (NaAlSi₃O₈), natrolite (Na₂Al₂Si₂O₇·2H₂O), and kieselguhr. With some of these typical catalysts, e.g., beryl, the following hydrocarbons (% vol.) were found in the product: C₁—16.1, C₂—38.5, C₃—32.7, C₄—12.7, and some C₅—C₁₀ derivs. The max. yield of liquid fraction of hydrocarbons was obtained in the instance of talc and kieselguhr catalysts. The effects of cryst. structure and traces of rare earth elements in the minerals are discussed in the light of their industrial utilization. A. P. Kuznetsov.

KAMZOLKIN, V.V., kandidat khimicheskikh nauk.

Producing higher aliphatic alcohols by direct oxidation of paraffin hydrocarbons. Khim. v shkole 11 no.2:26-27 Mr-Ap '56. (MIRA 9:7)
(Alcohols) (Hydrocarbons) (Oxidation)

AUTHOR: Bashkoriv, A.N., Kamzolkin, V.V., Sokova, K.M., and
Andreyeva, T.P. 65-4-2/12

TITLE: Method of determination of primary and secondary higher
alcohols of the fatty series in their mixtures. (Metod opre-
deleniya pervichnykh i vtorichnykh vyssikh spirtov zhirnogo
ryada v ikh smesyakh)

PERIODICAL: "Khimiya i Tekhnologiya Topliya i Masel" (Chemistry and
Technology of Fuels and Lubricants) 1957, No. 4, pp. 7-11 (U.S.S.R)

ABSTRACT: During studies of higher alcohols produced by a direct oxi-
dation of paraffinic hydrocarbons it was found difficult to
determine the content of primary and secondary alcohols, as
methods described in the literature (2, 3, 4) were found unsat-
isfactory when the number of carbon atoms in the molecules
exceeds eight. The method is based on some regularities in the
oxidation reaction of higher n-aliphatic alcohols with chromic
acid in glacial acetic acid. The accuracy of the method on av-
erage 5% (Table). There is one table and 6 references includ-
ing 3 Slavic.

Card 1/1

ASSOCIATION: Petroleum Institute Ac.Sc.U.S.S.R. (Institut Nefti
AN SSSR)

AVAILABLE:

KAMZOLKIN V.V.

BASHKIROV, A.N., doktor tekhnicheskikh nauk, professor; KAMZOLKIN, V.V.,
kandidat khimicheskikh nauk; LODZIK, S.A.

Technological elements of the production of higher fatty alcohols
by the direct oxidation of paraffinic hydrocarbons. Masl.-zhir.prom.
23 no.7:24-26 '57. (MLRA 10:8)

1. Institut nefti AN SSSR.
(Alcohols) (Hydrocarbons) (Oxidation)

KAMZOLKIN, V. V.

5(5) 12(4)

KAMZOLKIN, V. V.

Akademiya nauk SSSR, Institut nertsi

Trudy, t. 12 (Transactions of the Petroleum Institute, USSR. Academy of Sciences, Vol. 12) Moscow, Izdatel'nyy dom SSSR, 1978. 395 p. Extra slip inserted. 1,700 copies printed.

Ed.: S. B. Margulyash, Professor; Ed. of Publishing House: K. G. Myasov, Tech. Ed.: V. V. Golubina.

NOTE: The book is intended for scientists, engineers, and technicians in the petroleum industry.

CONTENTS: This collection of articles describes the results of studies on the chemistry and technology of petroleum and gas conducted in the laboratories of the Petroleum Institute, Academy of Sciences, USSR, in 1976 and 1977. A new section "Petrochemical Synthesis and Technology of Petroleum" has been included in the collection of articles. A list of investigations published by the scientists of the Institute in 1976 and 1977 and a list of dissertations for the Doctor's and Candidate's degrees presented in 1976 and 1977 at open sessions of the Academic Council of the Petroleum Institute, Academy of Sciences, USSR, are given. E. M. Izrael, P. V. Kuznetsov, I. A. Maslov, and V. V. Shchekin. Change in the Activity of Silicon Oil in the Chromatographic Separation of Hydrocarbons.

III. CARBONACEOUS AND CATALYTIC

Repin, Yu. B., A. M. Bakhitov, L. I. Zaslavskaya, and N. A. Orlov. Packed Bed Catalysts for the Synthesis of Higher Alcohols from Carbon Monoxide and Hydrogen. 200

Bakhitov, A. M., Ye. V. Kamzolkin, and Yu. B. Repin. Some Characteristics of the Decomposition of Carbon Monoxide into C and CO₂ in the Presence of Packed Bed Catalysts. 213

Legun, Yu. B., A. M. Bakhitov, B. M. Likhachev, K. G. Murty, and N. A. Orlov. Effect of Added Hydrocarbons on the Activity and Stability of Packed Bed Catalysts for the Synthesis from CO₂ and H₂. 228

Bakhitov, A. M., and P. I. Borok. Study of Conditions of Synthesis from Carbon Monoxide and Hydrogen in the Presence of Zinc Catalysts. 240

Mal'tin, S. A., A. Ya. Rozengolts, and V. V. Shchekin. Method of Kinetic Investigations of Continuous Gaseous Reactions. 246

Prokudin, L. P., A. Ya. Rozengolts, and V. V. Shchekin. Intradiusion Reactions in Catalytic Dehydration of Ethyl Alcohol. 253

Kozlovskaya, Z. V., and V. V. Shchekin. Adsorptive Properties of Aluminum Hydroxides and Aluminum Oxide. 261

Kozlovskaya, Z. V., and V. V. Shchekin. Activity and Structure of Aluminum Oxide and its Luminescent Properties. 267

Kozlovskaya, Z. V., and V. V. Shchekin. Anomalous Values of the Energy Constant of Fine-Pored Adsorbents. 272

Belitskiy, A. B., and V. M. Karicheva. Catalytic Addition of Hydrogen Chloride to Ethylene in Gaseous Phase. 276

IV. TECHNOLOGY OF PETROLEUM AND PETROCHEMICAL SYNTHESIS

Kamzolkin, V. V., A. M. Bakhitov, and M. Murty. Study of the Process of Continuous Oxidation of Paraffinic Hydrocarbons to Alcohols. 281

Kamzolkin, V. V., A. M. Bakhitov, and M. Murty. Investigation of the Effect of Nitric Acid and Boric Anhydride on the Liquid Phase Oxidation of Paraffinic Hydrocarbons. 290

Bakhitov, A. M., S. A. Lodyk, and V. V. Kamzolkin. Determination of the Content of Primary and Secondary Higher Alcohols by the Dehydration Method. 297

Krylov, Yu. B., V. K. Rubyshev, L. G. Ilyashov, E. A. Stepanov, and A. M. Bakhitov. Synthesis of Methyl Alcohol Containing the Radioactive Carbon Isotope, C¹⁴. 299

Kamzolkin, V. V., and L. V. Orlov. Manufacture of Acetonitrile by the Oxidation of Paraffinic Hydrocarbons with Ammonia in the Presence of Oxide Catalysts. 304

Belitskiy, A. B., (deceased), A. V. Ryzhenko, P. G. Anisimov, E. M. Belitskiy, low-temperature oxidative petroleum cracking. 321

Belitskiy, A. B., A. R. Brun-Tschow, Efficient Technology of Methane Conversion. 324

SOV5-58-6-3/13

AUTHORS: Bashkirov, A. N; Kamzolkin, V. V; Sekova, K. M. and Andreyeva, T. P.

TITLE: The Position of Hydroxyl Groups in Alcohols Prepared During the Liquid Phase Oxidation of n-Paraffin Hydrocarbons. (O.polezhenii gidroksil'nykh gruppy v spirtakh, poluchayemykh pri zhidkofaznom okislenii n-parafinovykh uglevodorodov).

PERIODICAL: Khimiya i Tekhnologiya Topliv i Masel, 1958, Nr.6. pp. 10 - 16. (USSR)

ABSTRACT: When investigating the position of the hydroxyl groups in the alcohol molecule, the authors used the method of oxidizing alcohols with potassium dichromate in a medium diluted with sulphuric acid (Ref.4). During the oxidation of primary alcohols, carboxylic acids, with the same number of C-atoms as contained in the initial alcohol, are obtained. During the oxidation of secondary alcohols, the C-C bonds are split at the hydroxyl groups, and carboxylic acids with a lower number of C atoms in the molecule are formed. Therefore, it is possible to determine the position of the hydroxyl groups according to the composition of the acids. Some side reactions take place when the process is carried out in sulphuric acid at increased temperatures. The authors investigated the

Card 1/3

SOV/65-58-6.3/13
The Position of Hydroxyl Groups in Alcohols Prepared During the Liquid
Phase Oxidation of n-Paraffin Hydrocarbons.

oxidation of individual aliphatic alcohols with varying positions of the hydroxyl group (4-tetradecanol and 7-hexadecanol) with subsequent identification of the acids. The method of F. Kraft (Ref.4) was slightly modified, and distillations were carried out according to the method described by L. K. Obukhova (Ref.5). The height of the rectification column was 40 cm and the diameter 1.4 cm. A mixture of hydrocarbons, from which the olefins and aromatic hydrocarbons had been separated, was used as carrier. The content of esters in the fractions was calculated on the basis of the ester number of the fraction. On the basis of the composition of the acids it was possible to conclude that oxidation of the alcohols occurs mainly at the hydroxyl groups. Discrepancies in the rule of Papev occur at increasing distances of the hydroxyl groups from the end hydrocarbon atom. The neutral oxygen-containing compounds (ketones), obtained during the oxidation, were subjected to second oxidation reaction. The total yield of acids is 96%. The investigated fractions of alcohols were concluded to be a mixture of isomers of secondary n-hexadecanols in which the isomers are contained in equal molar

Card 2/3

SOV/65-58-6-3/13

The Position of Hydroxyl Groups in Alcohols Prepared During the Liquid Phase Oxidation of n-Paraffin Hydrocarbons.

quantities. Experimental details on the oxidation of the individual alcohols are given. Tables 1 and 2 give the composition of oxidation products of alcohols and of their distillates; the distribution of acids is shown in Table 3. During experiments on defining the position of the hydroxyl groups in the alcohols, a fraction of alcohols boiling between 125.0 - 126.8, with an hydroxyl number of 229.5, was oxidized. The neutral oxygen-containing compounds were subjected to a second oxidation reaction. Results are given in Tables 4 and 5. These experiments showed that during the oxidation of n-paraffin hydrocarbons in the liquid phase, n-secondary alcohols are formed. The hydroxyl groups of these alcohols are situated at different C atoms of the molecule. It was also found that the reactivity of the secondary C atoms of molecules of higher n-paraffin hydrocarbons to oxygen is practically identical. There are 5 Tables and 8 References: 4 Soviet, 2 German, 1 English and 1 Dutch.

Card 3/3

ASSOCIATION: Petroleum Institute, AS USSR (Institut nefti AN SSSR)

KAMZOLKIN, V.V.

AUTHORS: Bashkirov, A. N., Kamzolkin, V. V.,
Sokova, K. M., Andreyeva, T. P.,

20-1-12/58

TITLE: On the Problem of the Oxidation Mechanism of Paraffinic Hydrocarbons in the Liquid Phase (K ~~vozniku~~ o mekhanizme zhidkofaznogo okisleniya parafinovykh ~~uglevodorodov~~)

PERIODICAL: Doklady AN SSSR, 1958, Vol. 118, Nr 1, pp. 149-152 (USSR)

ABSTRACT: This process is complicated and consists of a number of reactions taking place in parallel or successively. It is considered an established fact that this oxidation under mild conditions proceeds to water and carbonic acid through intermediate products of an incomplete oxidation (peroxides, alcohols, ketones, acids and others). A complicated mixture of oxygen-containing products develops. In an earlier paper (reference 1) the authors worked out the synthesis of higher alcohols of the aliphatic series by direct oxidation of paraffinic hydrocarbons in the presence of boric acid. The Shemism and the mechanism of individual stages has still to be determined. For this purpose the oxidation of a number of individual hydrocarbons was carried out and the composition of the alcohols produced was studied. A nitrogen-oxygen mixture (3,0 - 3,5% O₂) with addition of 5% boric acid (calculated on the initial hydrocarbon) under atmospheric pressure was

Card 1/3

On the Problem of the Oxidation Mechanism of Paraffinic Hydrocarbons in the Liquid Phase.

2-142/58

used for the oxidation. The temperature was 165-170°C and the duration was 4 hours. The hydrocarbons are characterized in table 1, the oxides ("oxydates"?) in table 2. From the latter follows that alcohols represent the main product (about 70%) of the oxidation. For determining their composition and structure they were isolated from the oxides. Their characteristics are recorded in table 3. From this is to be seen that the alcohols have hydroxyl numbers corresponding to tri-, tetra-, pent- and hexadecanole and consequently the same number of carbon atoms in the molecule as each of the corresponding initial hydrocarbons. In order to prove this alcohols were reconverted to hydrocarbons, in order to compare the properties of the latter with the initial hydrocarbons. The comparison of the two types of hydrocarbons showed their identity in the case of every individual alcohol. From this result the conclusion may be drawn that alcohols containing the same number of carbon atoms in the molecule as the initial hydrocarbons predominantly develop in the oxidation of n-paraffinic hydrocarbons by molecular oxygen in the liquid phase and under the conditions described. The molecule of the initial hydrocarbons is on the whole not destroyed. The determination of secondary alcohols meets with great difficulties

Card 2/3

On the Problem of the Oxidation Mechanism of Paraffinic Hydrocarbons in the Liquid Phase.

20-1-42/58

and does not lead to positive results. The methods known in publications (references 3,4) proved to be useless. In order to solve this problem the authors worked out a special method of the quantitative determination of primary and secondary alcohols. For this purpose alcohols were by means of chromic acid oxidized in the medium of glacial acetic acid. The accuracy of this method is about 5%. From the given results of analysis follows that predominantly secondary alcohols form in the reaction studied here (87,7 - 88,7 mol.%). The interaction of oxygen with the molecules of the paraffinic hydrocarbons of normal structure mainly takes place at the secondary carbon atoms. There are 4 tables, and 5 references, 3 of which are Slavic.

ASSOCIATION: Petroleum Institute AS USSR (Institut nefti Akademii nauk SSSR)
 PRESENTED: June 26, 1957, by A.V. Topchiyev, Academician
 SUBMITTED: June 26, 1957
 AVAILABLE: Library of Congress
 Card 3/3

The Synthesis of Ethanol From Carbon Dioxide and Hydrogen.

20-2-24/60

largest quantity of all. About 1% of the condensate was formed by acids, 90% of them acetic acid. Further the product desorbed from the activated charcoal was investigated (tables 3,4). It had been adsorbed from the outgoing gases during the synthesis. During the investigations some data on the chemism of the synthesis under review were obtained. An essential peculiarity of the process is the fact that it takes place in stages. At first carbon monoxide forms in quantities which do not exceed those in the water gas. Carbon monoxide represents the main source of the oxygen-containing compounds and of hydrocarbons. The concentration of carbon monoxide decreases with increasing concentration of water vapor in the reaction mixture. The synthesis comes to a standstill when the concentration of carbon monoxide sank to about 2%. The recirculation is capable of largely suppressing the formation of carbon monoxide. Table 5 gives some data on the products obtained in this connection. An approximate material balance shows a yield per 1 m³ gas mixture (CO₂ : H₂ = 1 : 3) in gram of: alcohols 92, hydrocarbons 81, water 345, carbon monoxide 31 and other oxygen-containing compounds 10. A lower speed of passage leads to an increase in the amount of acids. This may be in favor of the conception that the

Card 2/3

The Synthesis of Ethanol From Carbon Dioxide and Hydrogen.

20-2-24/60

main final product, ethanol, develops from acetic acid by reduction. This new synthesis of ethyl alcohol from carbon dioxide and hydrogen represents a step forward on the way of the development of the at present not yet numerous methods of converting carbon dioxide to valuable chemical products. There are 5 tables, 11 references, 4 of which are Slavic.

ASSOCIATION: Petroleum Institute, 'AS USSR (Institut nefti Akademii nauk SSSR)
PRESENTED: July 12, 1957, by A. V. Topchiyev, Academician
SUBMITTED: July 12, 1957
AVAILABLE: Library of Congress

Card 3/3

AUTHORS: Bashkirov, A. N., Kamzolkin, V. V. Sokova, K.M., 20-119-4-21/60
Andreyeva, T. P.

TITLE: The Composition of Alcohols Produced by Liquid Phase Oxidation of n-Paraffinic Hydrocarbons (O sostave spirtov, poluchayushchikh pri zhidkofaznom okislenii n-parafinovykh uglevodorodov)

PERIODICAL: Doklady Akademii Nauk SSSR, 1958, Vol. 119, Nr 4, pp. 705-707 (USSR)

ABSTRACT: The investigation of the chemical composition and the structure of higher aliphatic alcohols is very important for the explanation of their formation mechanism as well as for the selection of the right method for their rational exploitation. In the present paper the authors occupied themselves with the hydroxyl group in the alcohol molecule. They paid special attention to the method of oxidation of alcohols with sodium bichromate and with sulfuric acid. The weight ratios between alcohol and sodium bichromate were 1:3, the quantity of the diluted sulfuric acid and of alcohol 10:1. The oxidation was carried out at different temperatures. The existence of certain methyl ethers was found in single fractions. The

Card 1/3

The Composition of Alcohols Produced by Liquid Phase
Oxydation of n-Paraffinic Hydrocarbons

20-119-4-21/60

computation could be carried out also according to the
following formula:

$$X = \frac{A(M-1) - 107.9(100-A)}{14A} \cdot 100$$

whereby X denotes the content of acid C_n (mol.%); A denotes the content of silver in the existing silver salt (percentage by weight); M denotes the molecular weight of the acid C_{n+1} ; The numerical empirical data prove that this method of oxidation of alcohols can be used very well for the detection of the position of the hydroxyl group. The experimental results show furthermore that the alcohols obtained form a binary mixture of n-hexadecyl alcohols. The quantity of various alcohol molecules in this group is equal. Comprisingly was said that the alcohols produced by the oxidation of n-paraffinic hydrocarbons are mainly of secondary nature and represent a mixture of various isomeric substances.

The reactivity of the atoms of the molecules of higher paraffin hydrogens of normal structure does not display any considerable differences and is equal in comparison to oxygen. This is the condition for the production of isomeric substances

Card 2/3

The Composition of Alcohols Produced by Liquid Phase
Oxydation of n-Paraffinic Hydrocarbons

20-119-4-21/60

of secondary alcohols during the oxidation process of hydrocarbons in liquid state.
There are 1 table and 5 references, 4 of which are Soviet.

ASSOCIATION: Institut nefti Akademii nauk SSSR (Petroleum **Institute** of the AS USSR)

PRESENTED: December 14, 1957 by A. V. Topchiyev, Member, Academy of Sciences, USSR

SUBMITTED: December 14, 1957

Card 3/3

KADZOLKIN, V. V. , BASHKIROV, A. N.

"Synthesis of Higher Aliphatic Alcohols by the Method of Direct
Oxidation of Paraffinic Hydrocarbons."

Report, ^{the} submitted at the Fifth World Petroleum Congress, 30 May -
5 June 1959. New York.

Abstracts from USSR. Russian literature.

Oxidation of alcohols in acidic media; chemical theory (Oxidation of hydrocarbons in the liquid phase) Collection of Articles Moscow, 1958, 134 p. Russian only inserted. 2,200 copies printed.

M. I. M. Shumilov, Corresponding Member, Academy of Sciences USSR; M. I. of Publishing House: E. M. Izdatel'stvo, Moskva, M.I. I. P. Rus. 1958.

PURPOSE: This collection of articles is intended for scientists interested in hydrocarbon oxidation reactions, particularly for those specializing in petroleum fields.

CONTENTS: This collection of 35 articles represents the results of investigations over a period of several years on problems of hydrocarbon oxidation. The authors present their own theoretical and experimental data and also draw from current literature. No personalities are mentioned. The collection is the most of the articles.

Reaction of Alcohols with Peroxides, E. M. Shumilov, and V. P. Kozlovskaya (Institute of Chemistry of the USSR Academy of Sciences, Moscow, 1958). 159
Mechanism of Liquid-Phase Oxidation of Hydrocarbons in the Presence of Peroxides, E. M. Shumilov, and V. P. Kozlovskaya (Institute of Chemistry of the USSR Academy of Sciences, Moscow, 1958). 160
Oxidation of α -tridecane, α -pentadecane and α -heptadecane in the presence of borate acid is investigated. For the most part, secondary alcohols are formed. The purpose of the study was to establish a scientific basis for a method of preparing aliphatic alcohols by the direct oxidation of paraffin hydrocarbons developed by the Russian Institute.

Reaction of Alcohols with Peroxides, E. M. Shumilov, and V. P. Kozlovskaya (Institute of Chemistry of the USSR Academy of Sciences, Moscow, 1958). 160
Liquid-phase oxidation of hydrocarbons as a model reaction for the study of the mechanism of the oxidation of hydrocarbons in the presence of peroxides. The authors examine the mechanism of the oxidation of cyclohexane and discuss the various paths of cyclohexane formation. Liquid-phase oxidation of cyclohexane and methylcyclohexane in the presence of peroxides is investigated. The authors show that cyclohexane and methylcyclohexane decompose to form first a ketone, then an alcohol. Cyclohexane polymerization products were isolated and identified. The authors and trimers of cyclohexane hydroperoxides and their derivatives.

Reaction of Alcohols with Peroxides, E. M. Shumilov, and V. P. Kozlovskaya (Institute of Chemistry of the USSR Academy of Sciences, Moscow, 1958). 161
The authors conclude that the olefin oxidation process can be used to synthesize alcohols.

Reaction of Alcohols with Peroxides, E. M. Shumilov, and V. P. Kozlovskaya (Institute of Chemistry of the USSR Academy of Sciences, Moscow, 1958). 162
From their experiments on the oxidation of alcohols (hexanes), the authors conclude that the olefin oxidation process can be used to synthesize alcohols.

Reaction of Alcohols with Peroxides, E. M. Shumilov, and V. P. Kozlovskaya (Institute of Chemistry of the USSR Academy of Sciences, Moscow, 1958). 163
The authors examine the mechanism of the oxidation of cyclohexane and discuss the various paths of cyclohexane formation. Liquid-phase oxidation of cyclohexane and methylcyclohexane in the presence of peroxides is investigated. The authors show that cyclohexane and methylcyclohexane decompose to form first a ketone, then an alcohol. Cyclohexane polymerization products were isolated and identified. The authors and trimers of cyclohexane hydroperoxides and their derivatives.

Reaction of Alcohols with Peroxides, E. M. Shumilov, and V. P. Kozlovskaya (Institute of Chemistry of the USSR Academy of Sciences, Moscow, 1958). 164
The authors describe the separation and identification of hydroperoxides obtained in the oxidation of aliphatic-aromatic hydrocarbons, confirm the reactivity of various hydroperoxides, and report on the application of hydroperoxides as polymerization initiators.

Reaction of Alcohols with Peroxides, E. M. Shumilov, and V. P. Kozlovskaya (Institute of Chemistry of the USSR Academy of Sciences, Moscow, 1958). 165
The authors describe the separation and identification of hydroperoxides obtained in the oxidation of aliphatic-aromatic hydrocarbons, confirm the reactivity of various hydroperoxides, and report on the application of hydroperoxides as polymerization initiators.

Reaction of Alcohols with Peroxides, E. M. Shumilov, and V. P. Kozlovskaya (Institute of Chemistry of the USSR Academy of Sciences, Moscow, 1958). 166
The authors describe the separation and identification of hydroperoxides obtained in the oxidation of aliphatic-aromatic hydrocarbons, confirm the reactivity of various hydroperoxides, and report on the application of hydroperoxides as polymerization initiators.

Reaction of Alcohols with Peroxides, E. M. Shumilov, and V. P. Kozlovskaya (Institute of Chemistry of the USSR Academy of Sciences, Moscow, 1958). 167
The authors describe the separation and identification of hydroperoxides obtained in the oxidation of aliphatic-aromatic hydrocarbons, confirm the reactivity of various hydroperoxides, and report on the application of hydroperoxides as polymerization initiators.

Reaction of Alcohols with Peroxides, E. M. Shumilov, and V. P. Kozlovskaya (Institute of Chemistry of the USSR Academy of Sciences, Moscow, 1958). 168
The authors describe the separation and identification of hydroperoxides obtained in the oxidation of aliphatic-aromatic hydrocarbons, confirm the reactivity of various hydroperoxides, and report on the application of hydroperoxides as polymerization initiators.

5 (3)

AUTHORS: Kamzolkin, V. V., Bashkirov, A. N., SOV/20-126-6-38/67
Corresponding Member AS USSR,
Potarin, M. M.

TITLE: On the Synthesis of Higher Ketones by Means of the Oxidation of
Paraffin Hydrocarbons (O sinteze vysshikh ketonov metodom oki-
sleniya parafinovykh uglevodorodov)

PERIODICAL: Doklady Akademii nauk SSSR, 1959, Vol 126, Nr 6, pp 1282 - 1285
(USSR)

ABSTRACT: The formation of the carbonyl compounds takes place in the ox-
idation of the paraffin hydrocarbons at 120-160° much more ra-
pidly in the initial stage than the formation of the alcohols
and acids. On the basis of the concepts on the order of the for-
mation of oxygen containing compounds (see scheme, Refs 1,2) it
may be maintained that in this case ketones are formed mainly
directly from hydroperoxide. Because of this observation ex-
periments were carried out to determine whether a directed syn-
thesis of higher ketones by means of the oxidation of paraffin
hydrocarbons is possible in the liquid phase. For this purpose
the authors studied the action of the reaction conditions as
well as of some additions on the rate of oxidation and on the

Card 1/5

On the Synthesis of Higher Ketones by Means of the SOV/20-126-6-38/67
Oxidation of Paraffin Hydrocarbons

composition of some forming products. The fraction of synthetic hydrogenated paraffin hydrocarbons which boils out between 105 and 130° (1 torr) and which consists of carbons with 16 to 18 carbon atoms, was oxidized. The apparatus used and the method applied are described in reference 3. The temperature effects are shown by table 1. Their increase to 165° considerably intensifies the conversion of the initial hydrocarbons. A further increase to 185° remains practically without effect. The maximum yield of ketones was obtained at 120-140°. Also in the oxidation by a nitrogen oxygen mixture (3.5% O₂) the above regularities existed. By increasing this content to 21% O₂ the ketone

portion in the reaction products decreased (Table 2). At the same time, however, the degree of conversion of the initial hydrocarbons increased. It is possible that other oxidation conditions may be found under which high ketone yields are obtained. Such e.g. the reduction of the specific consumption of the oxidizing gas (oxygen) from 1000 l/kg .^h to 200 l/kg .^h leads to ketone yields of about 55% computed with respect to the reacted paraffin (Table 2). Table 3 shows the effect exercised by

Card 2/5

On the Synthesis of Higher Ketones by Means of the
Oxidation of Paraffin Hydrocarbons

SOV/20-126-6-38/67

the duration of oxidation on the composition of the oxide (3.5% O_2+N_2 at 140°). The oxidation intensity increases and the ketone yield decreases with the longer duration. The optimum duration is 3-4 hours in the oxidation with air at 140° . It may be seen from figure 1 that at 120° the oxidation is inhibited after a certain maximum degree of conversion (which depends on the reaction conditions) is attained. Later, the ketone and acid portion increases somewhat at the expense of the alcohol portions (Fig 1 and experiments Nr 1,2,4,5 in table 2). In this case an alcohol oxidation may take place. The oxidation process of the hydrocarbons concerned takes place according to the chain mechanism of the free radicals where a bimolecular decomposition of hydroperoxide leads to a branching of the chain, see scheme (Ref 4). The self-acceleration of the reaction at relatively low temperatures is due to the peroxide decomposition. The oxidation rate depends on the formation rate and the concentration of the free radicals in the reaction zone. The decomposition may take place also due to an interaction with the radical of reaction III according to reference 1. The mentioned inhibition is

Card 3/5

On the Synthesis of Higher Ketones by Means of the
Oxidation of Paraffin Hydrocarbons

SOV/20-126-6-38/67

probably due to a rapid reduction or stoppage of the formation of free radicals which are responsible of the reaction course. At higher temperatures no inhibition occurs and the oxidation takes place to a considerable degree of conversion (Fig 2). Alcohols, esters, acids and other oxygen containing compounds accumulate, however, in high quantities. Additions of KMnO_4 , FeSO_4 etc. could not bring about an increase of the ketone yield. The results obtained indicate that the formation of free radicals takes place mainly according to reaction (III) and not according to (I) and (II). Thus, it was proved that the higher aliphatic ketones may be produced by direct oxidation of paraffin hydrocarbons with yields of approximately 65 mol% of the transformed hydrocarbon and at a degree of conversion of 10-15 mol%. There are 2 figures, 3 tables, and 5 references, 3 of which are Soviet.

"ASSOCIATION: Institut neftekhimicheskogo sinteza Akademii nauk SSSR (Institute of Petrochemical Synthesis of the Academy of Sciences, USSR)
Card 4/5

5(2)

AUTHORS: Bashkirov, A. N., Corresponding Member, SOV/20-127-1-24/65
AS USSR, Potarin, M. M., Kamzolkin, V. V.

TITLE: The Synthesis of Higher Ketones by Liquid Phase Oxidation of
Secondary Alcohols (Sintez vysshikh ketonov metodom zhidko-
faznogo okisleniya vtorichnykh spirtov)

PERIODICAL: Doklady Akademii nauk SSSR, 1959; Vol 127, Nr 1, pp 93-96
(USSR)

ABSTRACT: The oxidation of the paraffin hydrocarbons in the liquid phase
is considered to be a process of various stages. The succession
of these stages is shown by a scheme (Refs 1-3). A considerable
number of carbonyl compounds are formed, as a rule, in the
production of synthetic sebacic acids by the oxidation of solid
paraffins. The oxidation products of the higher paraffin hydro-
carbons contain besides secondary alcohols (in the presence
of boric acid) always a certain quantity of ketones (Ref 5).
The carbonyl compounds (see Scheme) may be produced : a) by
the decomposition of hydroperoxide (Ref 6); b) by the oxidation
of the alcohols in the reaction zone. The authors attempted
to clarify the possibility of oxidizing the alcohols to ketones
under conditions similar to those of the oxidation of

Card 1/4

The Synthesis of Higher Ketones by Liquid Phase
Oxidation of Secondary Alcohols

SOV/20-127-1-24/65

hydrocarbons in the liquid phase, furthermore, the possibility of an orientated oxidation of secondary alcohols to ketones. For this purpose higher aliphatic alcohols (fraction boiling out within the range of from 125-170°, produced according to the method of reference 5) were used with a content of secondary alcohols of approximately 90 mole% (Ref 7). The apparatus and the method were already earlier described (Ref 5). The quantity of acids in the reaction products increases with rising temperature in the oxidation by means of an oxygen-nitrogen mixture (6.0-6.5 wt% O₂) during 4 hours at 120-180°, in contrast to that of ketones which is reduced from 63.6 to 55.3 mole% (Table 1). Carbonyl compounds apparently cannot be accumulated in greater quantities at higher temperatures since they are rapidly oxidized to acids. These acids occur either free or as esters. The transformation degree of the initial alcohols is increased by the increase of O₂-concentration in the reaction zone, the relative yield of ketones, however, is reduced (Table 2). The reaction rate depends as a rule to a considerable extent on the O₂-concentration in the oxidizing

Card 2/4

The Synthesis of Higher Ketones by Liquid Phase
Oxidation of Secondary Alcohols

SOV/20-127-1-24/65

gas. Its increase leads to greater yields of acids. The prolongation of the oxidation (Table 3, Fig 1) increases the transformation degree of the alcohols and the acid yield at the expense of the ketone yield. On the strength of the experimental results an oxidation method for alcohols could be chosen which guaranteed a ketone yield of 60 mole% in the case of a transformation degree of 40-45 mole% of alcohols; i.e.: temperature 165° , O_2 -content in the gas mixture

60-65 %, the specific consumption of the latter 1,000 l/kg^h, oxidation duration 3 hours. A higher yield of ketones may be obtained in the case of a lower transformation degree of the alcohols. Additions of Co-, Ni-, Mn-, Fe-, Cr-, Cu-, and Sn-salts are ineffective. Iron pentacarbonyl has a considerably inhibiting effect on the reaction. This proves its radical character. The data obtained are well in line with reference 8 with respect to the effect of boric acid and boric anhydride on the process mentioned in the title. There are 1 figure, 3 tables, and 8 Soviet references.

Card 3/4

The Synthesis of Higher Ketones by Liquid Phase
Oxidation of Secondary Alcohols

SOV/20-127-1-24/65

ASSOCIATION: Institut neftekhimicheskogo sinteza Akademii nauk SSSR
(Institute of Petrochemical Synthesis of the Academy
of Sciences, USSR)

SUBMITTED: March 16, 1959

Card 4/4

66172

~~5(3)~~ 5.3400

SOV/20-128-5-26/67

AUTHORS: Kamzolkin, V. V., Bashkirov, A. N., Corresponding Member, AS
USSR, Sokova, K. M., Andreyeva, T. P.

TITLE: On the Composition of Ketones Formed in Oxidation in the
Liquid Phase From n-Paraffin Hydrocarbons

PERIODICAL: Doklady Akademii nauk SSSR, 1959, Vol 128, Nr 5,
pp 956 - 959 (USSR)

ABSTRACT: Secondary alcohols with a normal structure are the main
products in the oxidation of n-paraffin hydrocarbons by mole-
cular oxygen in the liquid phase under mild conditions in the
presence of boric acid (Refs 1-3). The molecules of these
alcohols have the same number of carbon atoms as the initial
hydrocarbon. These alcohols form a mixture of all isomers
possible with regard to the position of the hydroxyl group.
Compounds with a carbonyl group, acids, and polyfunctional
compounds are formed as side products. The authors paid special
attention to the carbonyl compounds and used n-hexadecane
as initial product. Its oxidation took place in a device de-
scribed in reference 2. The oxidized substances enumerated in
table 1 were formed by the effect of a nitrogen-oxygen mixture

Card 1/3

66172

On the Composition of Ketones Formed in Oxidation
in the Liquid Phase From n-Paraffin Hydrocarbons

SOV/20-128-5-26/67

(with 3.5% O_2) at 165-170° and 5% boric acid within 4 hrs.

Among them were ~ 19 mol% compounds with a carbonyl group. The carbonyl compounds were separated from the oxidized substance for the purpose of determining their composition. About 40% of their total amount remained with the products which had reacted with boric acid. For this reason the authors assume that the afore-mentioned compounds containing carbonyl consist of ketones and polyfunctional compounds with a carbonyl group (mainly ketoalcohols). The distillate was chromatographically separated into paraffin hydrocarbons and compounds containing oxygen in order to determine the composition of the ketones. Table 2 shows the melting temperatures of semicarbazones. Comparison with them did not yield a final conclusion regarding the composition of the ketones investigated. Thus, they were oxidized with potassium bichromate in diluted H_2SO_4 (Ref 3).

Table 3 shows the results of the rectification of the methyl esters of the acids formed in the oxidized substance. Table 4 shows the weight- and per cent ratios of these acids. On account of the above results, the authors ascertain that mainly

Card 2/3

66172

On the Composition of Ketones Formed in Oxidation
in the Liquid Phase From n-Paraffin Hydrocarbons

SOV/20-128-5-26/67

ketones are represented among carbonyl compounds in the reaction mentioned. These ketones have the same chain length as the initial hydrocarbon. They form a mixture of all isomers theoretically possible with regard to the position of the carbonyl group. Polyfunctional compounds containing a carbonyl group are formed in addition to the ketones. There are 4 tables, and 4 references, 3 of which are Soviet.

ASSOCIATION: Institut neftekhimicheskogo sinteza Akademii nauk SSSR (Institute of Petroleum-chemical Synthesis of the Academy of Sciences, USSR)

SUBMITTED: June 26, 1959.

Card 3/3

BASHKIROV, A.N.; KAMZOLKIN, V.V.; SOKOVA, K.M.; ANDREYEVA, T.P.

Determination of primary and secondary higher alcohols of
the aliphatic series in their mixtures. Metod.anal.org.
soed.nefti,ikh smes. i proizv. no.1:170-177 '60. (MIRA 14:8)
(Alcohols) (Hydrocarbons)

KAMZOLKIN, V.V.; BASHKIROV, A.N.; SOKOVA, K.M.; ANDREYEVA, T.P.

Composition of oxygen-containing compounds formed in the
liquid phase oxidation of n-pentadecane by air. Trudy Inst.
nefti 14:65-75 '60. (MIRA 14:5)

(Pentadecane)

(Oxidation)

KAMZOLKIN, V.V.; BASHKIROV, A.N.; SOKOVA, K.M.; MARTYNES, M.; ANDREYEVA, T.P.

Transformations of higher aliphatic alcohols during their
liquid phase oxidation. Neftekhimiia 1 no.5:675-682 S-O '61.
(MIRA 15:2)

1. Institut neftekhimicheskogo sinteza AN SSSR.
(Alcohols)(Oxidation)

5.3400

69993

AUTHORS: Bashkirov, A. N., Corresponding Member of the AS USSR, Kamzolkin, V. V., Potarin, M. M., Kolovertnov, G. I. S/020/60/131/05/022/069 B011/B117

TITLE: Preparation of Higher Aliphatic Ketones by the Method of Dehydrogenation of Secondary Alcohols

PERIODICAL: Doklady Akademii nauk SSSR, 1960, Vol 131, Nr 5, pp 1067-1068 (USSR)

TEXT: The topic mentioned in the title has been studied using an industrial-type skeleton nickel catalyst. It was proved by the authors that it is possible to obtain a high yield (85% by weight), if the above-mentioned liquid-phase preparation method is used. The amount of the catalyst was 10% of the alcohol. Commercially produced C₁₆ - C₁₉ alcohols containing 6% of hydrocarbons were dehydrogenated. The reaction temperature was 185°, the residual pressure 33 torr. From the kinetic curves of the reaction it follows that the reaction proceeds rapidly in the liquid phase, and is practically completed within two hours. The conversion degree of the alcohols reaches 95 mole % (Fig 1). At first, a vigorous separation of hydrogen takes place, the iodine number of the product decreases, probably as a result of the hydrogenation of the unsaturated compounds in the alcohols used. Then, the iodine number is somewhat increased which is due to a side reaction involving the dehydration of the alcohols. It could be established

Card 1/2

6993

Preparation of Higher Aliphatic Ketones by the
Method of Dehydrogenation of Secondary Alcohols

S/020/60/131/05/022/069
BO11/B117

by chromatography on silica gel that the carbohydrate content was thereby increased from 6 to 10% by weight. The acid and ester content in the reaction products remains the same as the one in the alcohols used. The curves in figure 2 show that the dehydrogenation of the alcohols is accelerated by higher temperatures. Low pressure (33 torr) favors the reaction. The small quantity of unreacted alcohols was removed from the dehydrogenation product by esterification with boric acid. Substances not reacting with boric acid were distilled from the boric esters in vacuo (7 torr). The boiling-point range of the distillate was 115 to 120°. After removal of the hydrocarbons by means of chromatography on silica gel, a fraction of higher aliphatic ketones with d_4^{20} 0.8362, n_D^{20} 1.4446 and a carbonyl number of 202.0 was obtained. There are 2 figures and 5 references, 4 of which are Soviet.

ASSOCIATION: Institut neftekhimicheskogo sinteza Akademii nauk SSSR (Institute
of Petroleum-chemical Synthesis of the Academy of Sciences of the
USSR)

SUBMITTED: November 30, 1959

Card 2/2

Z/011/62/019/001/012/017
E073/E136

AUTHORS: Kamzolkin, V.V., and Bashkirov, A.N.

TITLE: Preparation of higher glycols by oxidation of paraffinic hydrocarbons in the liquid phase

PERIODICAL: Chemie a chemická technologie. Přehled technické a hospodářské literatury, v.19, no.1, 1962, 34, abstract Ch 62-473. (Neftekhimiya, v.1, no.3, 1961, 411-417)

TEXT: The experimental results indicate that during oxidation of paraffin hydrocarbons in the liquid phase in the presence of boric acid, secondary oxidation occurs of monofunctional oxygen compounds to multifunctional. During oxidation of esters by means of acetic acid the proportion of hydroxyl, carbonyl and multifunctional ester groups increases, which leads to the same conclusion. In the presence of acetic anhydride, the described method permits the preparation of glycol esters in addition to keto-alcohols.

1 figure, 4 tables, 7 references.

[Abstractor's note: Complete translation.]

Card 1/1

38689

S/510/60/014/000/003/006
D244/D307

5.3300

AUTHORS: Kamzolkin, V.V., Bashkirov, A.N., Sokova, K.M., and
Andreyeva, T.P.

TITLE: Composition of oxygen-containing compounds forming during
the liquid phase aerial oxidation of n-pentadecane

SOURCE: Akademiya nauk SSSR. Institut nefti. Trudy, v. 14, 1960,
Khimiya nefi, 65 - 75

TEXT: Results are presented of the study of the composition of the
products of oxidation of n-pentadecane in the presence of boric
acid. More oxygen was used in this work than previously (Bashkirov
A.N., Khimicheskaya nauka i promyshlennost', 1, no. 3, 272 (1956)).
The aim of the present investigation was to obtain additional data
on the oxidative conversions of hydrocarbons and on some intermedia-
te oxygen-containing compounds. It was found that the increase of O
in the oxidizing gas from 3.5 % to 21 % doubles the quantity of O -
containing compounds. At the same time the proportion of OH - con-
taining compounds decreases from 70 % to 50 % and COOH - containing
compounds increase from 12 % to 31 %. The amount of carbonyl com-
Card 1/2

Composition of oxygen-containing ...

S/510/60/014/000/003/006
D244/D307

pounds remains the same. Normal secondary alcohols are formed mainly with the OH group attached to different C atoms. The presence of poly-hydroxide compounds was established as well as the presence of a small quantity of primary alcohols containing less C atoms than the original hydrocarbon. Carbonyl compounds were found to be mainly ketones with 15 C atoms present in all possible isomeric forms. Acids not soluble in H₂O contained different molecules of different molecular size, with the highest percentage of acids lying in the C₈ - C₁₀ range. Compounds with different functional groups in the same molecule were also found. The results obtained confirm that the oxidative conversion of hydrocarbons is a single stage process. There are 9 tables. ✓

Card 2/2

KAMZOLKIN, V.V.; BASHKIROV, A.N.; KHOTIMSKAYA, M.I.; GROZHAN, M.M.;
YEZHENKINA, G.M.

Synthesis of aliphatic C₆ - C₁₀ alcohols by the liquid phase
oxidation of paraffins under pressure. Neftekhimiia 1 no.2:
244-254 Mr-Apr '61. (MIRA 15:2)

1. Institut neftekhimicheskogo sinteza AN SSSR.
(Alcohols) (Oxidation) (Paraffins)

KAMZOLKIN, V.V.; BASHKIROV, A.N.; LODZIK, S.A.

Composition of alcohols obtained by the direct oxidation of
paraffins under industrial conditions. Neftekhimiia 1 no.2:
260-266 Mr-Ap '61. (MIRA 15:2)

(Alcohols)
(Paraffins)

KAMZOLKIN, V.V.; BASHKIROV, A.N.; LODZIK, S.A.

Possibilities for obtaining higher glycols by liquid-phase
oxidation of paraffin hydrocarbons. Neftekhimia 1 no.3:
411-417 My-Je '61. (MIRA 16:11)

1. Institut neftekhimicheskogo sinteza AN SSSR.

BASHKIROV, A.N.; KAMZOLKIN, V.V.; SOKOVA, K.M.; ANDREYEVA, T.P.;
KORNEVA, V.V.; ZAKHARKIN, L.I.

Synthesis of cyclododecanol by the liquid-phase oxidation
of cyclododecane. Neftekhimiia 1 no.4:527-534 J1-Ag '61.
(MIRA 16:11)

1. Institut neftekhimicheskogo sinteza AN SSSR i Institut
elementoorganicheskikh soedineniy AN SSSR.

ZAKHARKIN, L.I.; KORNEVA, V.V.; KAMZOLKIN, V.V.; SOKOVA, K.M.;
ANDREYEVA, T.P.; BASHKIROV, A.N.

Preparation of ω -dodecalactam from 1,5,9-cyclododecatriene.
Neftekhimiya 2 no.1:106-109 Ja-F '62. (MIRA 15:5)

1. Institut elementoorganicheskikh soedineniy AN SSSR.
(Lactams) (Cyclododecatriene)

5

ZAKHARKIN, I.I., KOREVA, V.V., KURITSBERAYA, G.M., BASHKIROVA, A.N.,
KAMZOLKIN, V.V., SONOVA, K.M.

New monomer for the production of the synthetic fiber dode-Kalaktan.

Report to be submitted for the 12th Conference on high molecular weight compounds
devoted to monomers, Baku, 3-7 April 62

KOLESNIKOVA, L.P.; KAMZOLKIN, V.V.; KHOTIMSKAYA, M.I.; Primala
uchastiye: STAROBINETS, L.L.

Use of gas chromatography in the study of isomeric composition
of alcohols. Neftekhimiia 2 no.3:355-358 My-Je '62.

(MIRA 15:8)

1. Institut neftekhimicheskogo sinteza AN SSSR.
(Alcohols) (Gas chromatography)

KAMZOLKIN, V.V.; BASHKIROV, A.N.; KAMZOLKINA, Ye.V.; LODZIK, S.A.

Certain laws governing the liquid-phase oxidation of olefins.
Neftekhimiia 2 no.5:750-755 S-O '62. (MIRA 16:1)

1. Institut neftekhimicheskogo sinteza AN SSSR.
(Olefins) (Oxidation)

KAMZOLKIN, V.V.

2

SOKOLOVA, S.M., KAMZOLKIN, V.V., ANDREYEVA, T.P.

Obtaining cyclododecanol by liquid-phase oxidation of cyclododecane

Report to be submitted for the Sixth World Petroleum Congress,
Frankfurt, 16-26 June 63

KRYUKOV, Yu.B.; SMIRNOVA, R.M.; SELEZNEV, V.A.; KAMZOLKIN, V.V.;
BASHKIROV, A.N.

Intermediate stages in the liquid phase oxidation of secondary
alcohols to ketones. Neftekhimiia 3 no.2:238-245 Mr-Apr '63.
(MIRA 16:5)

1. Institut neftekhimicheskogo sinteza AN SSSR imeni A.V.Topchiyeva.
(Alcohols) (Oxidation) (Ketones)

KAMZOLKIN, V.V.; BASHKIROV, A.N.; SOKOVA, K.M.; ANDREYEVA, T.P.

By-products of the liquid-phase oxidation of cyclododecane with
molecular oxygen in the presence of boric acid. *Neftekhimiya*
4 no.1:96-99 Ja-F'64 (MIRA 17:6)

1. Institut neftekhimicheskogo sinteza AN SSSR imeni A.V.
Topchiyeva.

BASHKIROV, A.N.; KAMZOLKIN, V.V.; POTARIN, E.M.

Obtaining higher ketones by the dehydrogenation of secondary
alcohols over copper-chromium and nichel-chromium catalysts.
Neftekhimiia 4 no.2:298-300 Mr-Apr'64 (MIRA 17:8)

1. Institut neftekhimicheskogo sinteza AN SSSR imeni A.V. Topchi-
yeva.

PONOMARENKO, A.T.; KAGAN, Yu.B.; KAMZOLKIN, V.V.

Device for measuring gas consumption under high pressure. Khim.
i tekhn. topl. i masel 9 no.2:48-50 F '64. (MIRA 17:4)

1. Institut neftekhimicheskogo sinteza AN SSSR.

NOVAK, F.I.; KAMZOLKIN, V.V.; BASHKIROV, A.N.

Catalytic activity of the natural silicates of minerals in the
synthesis of hydrocarbons from carbon monoxide and hydrogen.
Neftekhimia 4 no.3:447-451 My-Je '64. (MIRA 18:2)

1. Institut neftekhimicheskogo sinteza AN SSSR im. A.V.Topchiyeva.

KAMZOLKIN, V.V.; KRYUKOV, Yu.B.; KAGAN, Yu.B.

Prospective trend of the petroleum chemistry. Vest. AN SSSR 34
no.11:60-65 N '64. (MIRA 17:12)

1. Institut neftekhimicheskogo sinteza im. A.V. Topchiyeva AN SSSR.

KAMZOLKINA, N. B.: Master Med Sci (diss) -- "The antigen structure of the New-castle dysentery microbe and its serological and immunological interrelationship with microbes of the Flexner sub-type". Moscow, 1958. 15 pp (First Moscow Order of Lenin Med Inst im I. M. Sechenov), 200 copies (KL, No 7, 1959, 129)

KEMZOLKINA, N.B.; LUKASHOVA, N.I.; ZAKHAROVA, N.S.; BORISOVA, L.V.

Use of cellular cultures for the determination of antitoxin
content in antidiphtheria sera. Zhur. mikrobiol., epid. i immun.
42 no.11:122-123 N '65. (MIRA 18:12)

1. Submitted April 14, 1965.

RPSHTEYN-LITVAK, R.V.; DMITRIYEVA-RAVIKOVICH, Ye.M.; D'YAKOVA, Ye.I.;
KAMENSKAYA, I.N.; VIL'SHANSKAYA, F.L.; KAMZOLKINA, N.B.

Theoretical bases of dysenterial immunity. Zhur. mikrobiol. epid. i
immun. 32 no.6:18-25 Je '61. (MIRA 15:5)

1. Iz Moskovskogo instituta epidemiologii, mikrobiologii i gigiyeny.
(DYSENTERY) (IMMUNITY)

KAMZOLKINA, Ye. V.

USSR/Chemistry - Synthetic Liquid Fuels Jul/Aug 52

"The Role of Oxygen-Containing Compounds in the Synthesis of Hydrocarbons From Carbon Monoxide and Hydrogen," Yu. B. Kagan, Yu. B. Kryukov, Ye. V. Kamzolkina, A. N. Bashkirov, Petroleum Inst. Acad Sci USSR "Iz Ak Nauk SSSR, Otdel Khim Nauk" No 4, pp 649-657, 1952

Article states that results of the expts described show that alcs cannot be regarded as intermediate products in the synthesis of hydrocarbons, and that iron catalysts, under the conditions of hydrocarbon synthesis, accelerate the oxidation of alcs and aldehydes. Advances hypotheses explaining the formation of oxygen-contg compds (by-products of hydrocarbons).

PA 229T16

KAMZOLKIN, Ye. U.

5(5); 12(4) PAGE 1 BOOK DEPOSITED 06/7/2021

Akademys nauk SSSR, Institut verti

Prely, 6, 12 (Transactions of the Petroleum Institute, USSR, Academy of Sciences, Vol. 12) Moscow, Izdat. AN SSSR, 1958. 595 p. First's slip inserted, 1,700 copies printed.

MA.1 E. E. Sergiyenko, Professor, Ed. of Publishing House: E. O. Kuznetsov, Mosk. Ed., T. T. Dolobova.

PURPOSE: The book is intended for scientists, engineers, and technicians in the petroleum industry.

CONTENTS: This collection of articles describes the results of studies on the activity and mechanism of petroleum and gas conducted in the laboratories of the Petroleum Institute, Academy of Sciences, USSR, in the field of catalysis. The book contains a large number of articles, a list of references, has been included in the collection of articles. A list of investigations published by the associates of the Institute in 1956 and 1957 and a list of dissertations for the Doctor's and Candidate's degrees presented in 1956 and 1957 at open sessions of the Academic Council of the Petroleum Institute, Academy of Sciences, USSR, are given. General of the Petroleum Institute, I. A. Musyev, and V. V. Shchekin. Change in the Activity of Nickel Oil in the Chromatographic Separation of Hydrocarbons

III. CATALYSIS AND CHEMISTRY

- 200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000

- 276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000

KAMZOLKINA, Ye. V., Candidate Chem Sci (diss) -- "A study of the decomposition reactions of carbon monoxide on melted iron catalysts for synthesis from carbon monoxide and hydrogen". Moscow, 1959. 13 pp (Acad Sci USSR, Inst of Petroleum-Chem Synthesis), 120 copies (KL, No 23, 1959, 161)

5(2)

SOV/156-59-1-42/54

AUTHORS:

Bashkirev, A. N., Kamzolkina, Ye. V., Kagan, Yu. B.

TITLE:

On Catalysts of the Reactions of the Decomposition of Carbon Monoxide (O katalizatorakh reaktsiy razlozheniya okisi ugloroda)

PERIODICAL:

Nauchnyye doklady vysshey shkoly. Khimiya i khimicheskaya tekhnologiya, 1959, Nr 1, pp 162 - 165 (USSR)

ABSTRACT:

For the systematic investigation of the influence of catalysts on the reactions: 1) $2 \text{Fe} + 2 \text{CO} \rightarrow \text{Fe}_2\text{C} + \text{CO}_2$, and 2) $2\text{CO} \rightarrow \text{C} + \text{CO}_2$, melted iron catalysts with activating additives (Al_2O_3 , SiO_2 , K_2O , Cr_2O_3 , B_2O_3 , MoO_3 , V_2O_5 , MgO , MnO , and combinations thereof), as employed in the synthesis $\text{Co}+\text{H}$, were investigated. The results are listed in a table. Carbide formation (reaction 1) and reaction 2 were accelerated by Al_2O_3 and V_2O_5 . K_2O accelerates only reaction 1. MoO_3 is ineffective, Cr_2O_3 , FeCr , and in particular SiO_2 and B_2O_3 exercise inhibitory effects. With several additives to the catalyst, the properties have, as a rule, a cumulative effect.

Card 1/2

On Catalysts of the Reactions of the Decomposition of
Carbon Monoxide

SCV/156-59-1-42/54

Only FeCr, which, when added exclusively, inhibits the reaction, increases the reaction velocity in the presence of other additives. Thus an iron catalyst with kaolin (as an SiO_2 vehicle), K_2O and FeCr showed the highest reactivity.

In this case, even B_2O_3 inhibits only reaction 2. A further test series concerned industrial iron catalysts, which were also given additives. The table of the results shows that also in this case the same rules apply. There are 2 tables and 1 Soviet reference.

ASSOCIATION: Kafedra neftekhimicheskogo sinteza i iskusstvennogo zhidkogo topliva Moskovskogo instituta tonkoy khimicheskoy tekhnologii im. M. V. Lomonosova (Chair of Petroleum-chemical Synthesis and Artificial Liquid Fuels of the Moscow Institute of Fine Chemical Technology imeni M. V. Lomonosov)

SUBMITTED: October 6, 1958

Card 2/2

S/195/60/001/003/008/013
B013/B058

AUTHORS: Kagan, Yu. B. Bashkirov, A. N., Kamzolkina, Ye. V.,
Loktev, S. M.

TITLE: On the Activation Process of Molten Iron Catalysts for
for CO and H₂ Synthesis Under the Effect of the Reaction
Mixture

PERIODICAL: Kinetika i kataliz, 1960, Vol. 1, No. 3, pp. 393 - 400

TEXT: The activation of molten iron catalysts for the hydrocarbon
synthesis from CO and H₂ under the effect of the reaction mixture was
studied in this paper. The following catalysts were used:

- 1) $100\text{Fe}_3\text{O}_4 + 6\text{Al}_2\text{O}_3 + 4.2\text{SiO}_2 + 1.2\text{K}_2\text{O} + 0.3\text{Cr}$;
- 2) $100\text{Fe}_3\text{O}_4 + 6\text{Al}_2\text{O}_3 + 4.2\text{SiO}_2 + 1.2\text{K}_2\text{O} + 0.5\text{V}$;
- 3) $100\text{Fe}_3\text{O}_4 + 6\text{Al}_2\text{O}_3 + 4.2\text{SiO}_2 + 1.2\text{K}_2\text{O} + 1.0\text{B}_2\text{O}_3$. They were reduced

Card 1/4

On the Activation Process of Molten Iron
Catalysts for CO and H₂ Synthesis Under the
Effect of the Reaction Mixture

S/195/60/001/003/008/013
B013/B058

within 1.5 hrs in hydrogen current at 1000°C. The study was made in a highpressure apparatus (Ref. 4) in the laboratory. For each of the catalysts studied, the lowest temperatures and pressures were initially chosen, at which, over the freshly and reduced catalysts (in comparable time intervals), a high degree of transformation of the carbon monoxide (84 to 86%) entering at a volume rate of the initial gas (CO and H₂ 1 : 1) of ~1500 h⁻¹ was obtained. The catalysts were gradually activated under these conditions. The duration of the tests varied. The tests of catalysts of equal composition were conducted under the same conditions and in the same reaction vessel. The indices of the synthesis were well reproducible. The results determined could therefore also be compared with each other. The samples were hydrogenated after termination of the synthesis test. Subsequently, the hydrogenated samples were treated with CO at atmospheric pressure, a volume rate of 800 h⁻¹ and temperatures by 10° higher than at the end of the synthesis test, with carbide being formed. The studies

Card 2/4

On the Activation Process of Molten Iron
Catalysts for CO and H₂ Synthesis Under the
Effect of the Reaction Mixture

S/195/60/001/003/008/013
B013/B058

produced the following results: the previously (Refs. 1, 2) made statement that the iron catalysts molten at high temperatures (1000°C) immediately after reduction, are inactive in the synthesis of CO and H₂ and obtain

activity only during the course of this synthesis, was confirmed. Activation also continues after reaching activity, which warrants a practically complete transformation of the initial carbon monoxide. This process is not terminated until 3 to 4 days after conduction of the synthesis. Simultaneously with the activation of the catalysts under the effect of the reaction mixture, their reactivity with respect to carbide formation is also increased. Those catalysts which have reached equal activity in consequence of the CO + H₂ synthesis, have also a similar reactivity with regard to carbide formation. The activation of the catalysts during the synthesis is accompanied by an increase of their activity during CO decomposition under formation of elementary carbon. The conditions under which the activation of catalysts occur (pressure, temperature, CO₂ content of the gas) have a noticeable effect on their properties. Of the methods investigated of

Card 3/4

On the Activation Process of Molten Iron
Catalysts for CO and H₂ Synthesis Under the
Effect of the Reaction Mixture

S/195/60/001/003/008/013
B013/B058

the activation of the catalyst for the synthesis, its treatment at the
synthesis temperature is suited best. In this case the catalysts get speci-
ally active, but simultaneously show a lower activity with regard to CO
decomposition. There are 3 figures, 4 tables, and 6 Soviet references.

ASSOCIATION: Institut neftekhimicheskogo sinteza AN SSSR
(Institute of Petrochemical Synthesis AS USSR)

SUBMITTED: January 19, 1960

Card 4/4

KAMZOLKIN, V.V.; BASHKIROV, A.N.; KAMZOLKINA, Ye.V.; LODZIK, S.A.

Certain laws governing the liquid-phase oxidation of olefins.
Neftekhimiia 2 no.5:750-755 S-O '62. (MIRA 16:1)

1. Institut neftekhimicheskogo sinteza AN SSSR.
(Olefins) (Oxidation)

BRODSKIY, Matvey Abramovich; KAMZOLOV, Aleksey Pavlovich; KASPEROVICH, N.S.,
inzh., red.; UVAROVA, A.F., tekhn. red.

[Catalog of spare parts for agricultural tractors] Katalog zapasnykh
chastei sel'skokhoziaistvennykh traktorov. Moskva, Gos. nauchno-
tekhn. izd-vo mashinostroit. lit-ry, 1961. (MIRA 14:8)

(Tractors--Catalogs)

KAMZOLOV, N. F.

Self - determination, National

National and colonial problem in the decisions of the Prague Conference. Vest. Mosk.
un. 7 no. 4, 1952.

9. Monthly List of Russian Accessions, Library of Congress, August, 1952 ~~1953~~. Unclassified.

LEONOV, O.B.; kand.tekhn.nauk, dotsent; KAMZOLOV, Ye.P., aspirant

Investigating film carburation. Izv.vys.ucheb.zav.; mashinostr.
no.1:116-122 '61. (MIRA 14:4)

1. Moskovskoye vyssheye tekhnicheskoye uchilishche imeni Baumana.
(Diesel engines---Testing)

KAMZOLOV, Ye.P., aspirant

Investigating the process of ignition and combustion of fuel
evaporating from a heated surface. Izv.vys.ucheb.zav.; mashinostr.
no.4:124-132 '61. (MIRA 14:6)

1. Moskovskoye vyssheye tekhnicheskoye uchilishche imeni Baumana.
(Combustion)

32240

S/145/61/000/004/004/008

D221/D301

11.7100
AUTHOR:

Kamzolov, Ye.P., Aspirant

TITLE:

Investigating the process of fuel ignition and combustion process during its evaporation from a heated surface

PERIODICAL: Izvestiya vysshikh uchebnykh zavedeniy. Mashinostroyeniye, no. 4, 1961, 124 - 132

TEXT: The investigation was carried out on a M4 (MCh) 10.5/13 engine with a cylindrical turbulent combustion chamber. One of the covers of the detachable chamber held a piezo-quartz transducer for measuring pressure, whereas the other cover contained a heat resisting optical glass JK-5 (LK-5). This permitted high speed photography of the process. The required temperature was established by electric heaters, controlled by two thermocouples. The fuel system incorporated two nozzles - a working and a supplementary unit. The cyclic operation was ensured by automatic blocking of the auxiliary nozzle. The incidence angle of the flame was varied by ex-

Card 1/5

32240
S/145/61/000/004/004/008
D221/D301

Investigating the process of ...

change of atomizers. The angles between the axes of nozzle apertures and the atomizer α , and the angle of the axis of the flame and the tangent to the surface of the combustion chamber β , (called the injection angle), as well as the free length of flame before colliding with the wall l_f are illustrated in Fig. 2. The instruments used allowed simultaneous measurement (and photography) of processes, recording of pressure in the combustion chamber p_z , injection pressure p_i , lift of the needle in the working nozzle, and the marking of the upper dead center as well as the time. The experiments were carried out with a diesel fuel АЛ(DL) and automobile gasoline А-70 (A-70). The plot of the temperature effect reveals a slight increase of p_z and of the rate of pressure incre-

ment, $\frac{\Delta p}{\Delta \varphi}$ with higher heating temperature, t_h . This may be due to faster spread of the flame. In addition, reduced lagging of the ignition is partly compensated by faster evaporation. This ensures a smooth engine operation within a wide range of surface temperature.

Card 2/5

32240

S/145/61/000/004/004/008

D221/D301

Investigating the process of ...

res t_h . A low t_h is uneconomical. The two-center ignition with an atomizer of $\alpha = 70^\circ$ at $t_h = 350^\circ \text{C}$ has changed to a single center ignition when this temperature went down. The diagram concerning the effect of injection angle demonstrates that in the case of a single nozzle, low α results in higher p_z and $\frac{\Delta p}{\Delta \varphi}$. Reduction of α produces a drop of ignition lag τ , higher speed of flame spreading, and shorter combustion time. This increases the importance of mixing when α decreases. Rational design of the atomizer is determined by $\frac{\Delta p}{\Delta \varphi}$ and the economy, which depends on τ_c . A two-nozzle unit ensures a better start, but has design disadvantages. The increase of p_z and of $\frac{\Delta p}{\Delta \varphi}$ with the angular velocity is explained by higher speed of air charge, injection and flame spread. Greater values of the advance angle θ produce a fuel injection at lower temperatures. Therefore, the chemical and physical preparation of the fuel for the self-ignition is retarded. The investigations permit the fol-

Card 3/5

32240

S/145/61/000/004/004/008

D221/D301

Investigating the process of ...

lowing conclusions: The change of surface temperature from 250 to 350°C has little effect on the process. Further increase of heating may cause undesirable abruptness in the operation. The angle β is the main design parameter. The alteration of speed is reflected on the course of ignition and combustion processes. The optimum advance angle of injection in the case of film mixing is somewhat greater than for volumetric mixing. Excess in this quantity increases the rigidity of operation. There are 7 figures. ✓

ASSOCIATION: MVTU im. N.E. Bauman (MVTU im. N.E. Bauman)


SUBMITTED: October 31, 1960

Card 4/5

Investigating the process of ...

S/145/61/000/004/004/008
D221/D301

Fig. 2.



Card 5/5

KAMZOLOVA, K. P.

AKOVBYAN, A.A., professor; KAMZOLOVA, K.P., dotsent (Tashkent)

Methods of teaching dermatovenereology in medical institutes. Vest.
derm. i ven. 31 no.4:29-30 J1-Ag '57. (MIRA 10:11)

(DERMATOLOGY, educ.

dermato-venereol., teaching methods in Russia)

(VENEREAL DISEASES

same)

KAMZOLOVA, K.P.; TERNOVENKO, K.M.

Armais Aristogesovich Akovbian. Med. zhur. Uzb. no.6:76-77 Je '60.
(MIRA 15:2)

(AKOVBIAN, ARMAIS ARISTOGESOVICH, 1900-)

AKOVBYAN, A.A., prof.; KAMZOLOVA, K.P., dotsent

Treatment vitiligo with Soviet psoralen. Vest.derm.i ven. no.5:
33-38 '61. (MIRA 14:12)

1. Iz kafedry kozhnykh i venericheskikh bolezney (zav. - prof.
A.A. Akovbyan) Tashkentskogo meditsinskogo instituta.
(SKIN---DISEASES) (PSORALEN)

KUCHEROVA, N.F.; ZHUKOVA, I.G.; KAMZOLOVA, N.N.; PETRUCHENKO, M.I.;
SHARKOVA, N.M.; KOCHETKOV, N.K.

Indole derivatives. Part 8: 9-Acyl-1,2,3,4,4a,9a-hexahydro-8-carbolines. Zhur.ob.khim. 31 no.3:930-936 Mr '61, (MIRA 14:3)

1. Nauchno-issledovatel'skiy institut farmakologii i khimioterapii.
(Pyridindole)

DREYMAN, E.Ya.; DMITRIYEVA, V.A.; KAMZOLOVA, S.G.; SHABAROVA, Z.A.;
PROKOF'YEV, M.A.

Synthesis of o-aminoacyl derivatives of adenosine and study of
their properties. Zhur.ob. chim. 31 no.12:3899-3905 D '61.
(MIRA 15:2)

1. Moskovskiy gosudarstvennyy universitet imeni M.V.Lomonosova.
(Adenosine)

NRAP0033074

SOURCE CODE: UR/0218/66/031/005/0910/0917

AUTHOR: Shemyakin, M. P.; Bass, I. A.; Kamzolova, S. G.; Gorlenko, Zh. M.; Astaurova, O. B.; Khesin, R. B.

ORG: Order of Lenin Atomic Energy Institute im. I. V. Kurchatov, Moscow (Ordena Lenina institut atomnoy energii)

TITLE: Specificity of RNA synthesis in phage infection

SOURCE: Biokhimiya, v. 31, no. 5, 1966, 910-917

TOPIC TAGS: RNA, RNA synthesis, infective disease, bacteriophage, biochemistry, biosynthesis, ~~E. coli~~, ~~T2 phage~~, polymerase, ~~RNA~~

ABSTRACT: The specificity of RNA synthesis in different phases of T2 bacteriophage infections of *E. Coli* B and in an *in vitro* RNA polymerase system was investigated using labeled RNA. In early and late infectious stages, mRNA is synthesized largely on different regions of the T2 phage chromosome. Results of *in vitro* experiments show that RNA polymerase synthesizes RNA on the same regions of purified T2 phage which are active in intact cells during early stages of infection. Orig. art. has: 3 fig. and 1 table [LP]

SUB CODE: 06/ SUBM DATE: 15Nov65/ ORIG REF: 004/ OTH REF: 014 [WA-50; CBE No. 14]
Card 1/1 UDC:547.963.3

VASILENKO, S.K.; KAMZOLOVA, S.G.; KNORRE, D.G.

Direct spectrophotometric method for the quantitative determination of the nucleotide composition of ribonucleic acids. Biokhimiia 27 no.1:142-148 Ja-F '62.
(MIRA 15:5)

1. Institute of Organic Chemistry, the Siberia Branch of Academy of Sciences of the U.S.S.R., Novosibirsk.
(NUCLEOTIDES) (SPECTROPHOTOMETRY) (NUCLEIC ACIDS)

1ST AND 2ND COLUMNS										3RD AND 4TH COLUMNS									
COMMON ELEMENTS										COMMON VARIABLES INDEX									
KAMENKOVA, L.										PROCESSING AND PROPERTIES INDEX									
BC										a-1									
<p>Viscometric investigations of structure formation in ferric hydroxide sols. III. Destruction of ferric hydroxide sols on dilution. G. FUCHS and L. KAMENKOVA (Acta Physicochim. U.R.S.S., 1955, 3, 127-134). The destruction of young and aged conc. $Fe(OH)_3$ sols and gels on dilution has been studied. The effect of adding $FeCl_3$ was also examined. The viscosity data indicate the existence of structures of different ages and solidity in the sol particles, which may be termed primary and secondary structures. The mechanism of destruction consists in the breakdown of secondary structures and displacement of the adsorption equilibrium with dilution. A. J. M.</p>																			
ASB-SLA METALLURGICAL LITERATURE CLASSIFICATION																			
SOURCE SYMBOLISM										SOURCE SYMBOLISM									
SOURCE NO.										SOURCE NO.									
SOURCE NO.										SOURCE NO.									

KANZOLOVA, Z.P.

~~MINERALOGY OF DRY CLAYS OF THE ANGREN REGION~~

Mineralogy of "dry" clays of the Angren region. Zap.Uz.otd.Vses.min.ob-
va no.2:104-108 '51. (MLRA 6:6)

(Angren Region--Clay)

S/129/61/000/004/010/012
E073/E535

AUTHORS: Shreyber, G. K., Candidate of Technical Sciences and
Kan, A., Engineer

TITLE: Residual Stresses During Surface Hardening of Steel

PERIODICAL: Metallovedeniye i termicheskaya obrabotka metallov,
1961, No.4, pp.44-49

TEXT: The authors attempted to determine analytically the magnitude and the character of residual stresses which arise during surface hardening in various zones of a cylindrical steel specimen. The residual thermal stresses in a solid cylindrical specimen can be determined by means of the following approximate formulae.

1. Hardened zone:

$$\sigma_{ps} = \frac{\alpha TE}{4(1-\mu)} \left(1 + \frac{R_n^2}{R^2} - \frac{r^2}{R^2} - \frac{k_n^2}{r^2} \right), \quad (1)$$

$$\sigma_{rs} = -\frac{\alpha TE}{4(1-\mu)} \cdot \left(3 \frac{r^2}{R^2} - \frac{R_n^2}{r^2} - \frac{R_n^2}{k^2} - 1 \right), \quad (2)$$

Card 1/10

Residual Stresses During Surface....

S/129/61/000/004/010/012
E073/E535

$$\sigma_{\theta\theta} = -\frac{\alpha TE}{2(1-\mu)} \left(2 \frac{r^2}{R^2} - \frac{R_n^2}{R^2} - 1 \right). \quad (3)$$

2. Transition and Unhardened zones:

$$\sigma_{P_n} = -\frac{\alpha TE}{4(1-\mu)} \times \frac{R^2 - R_n^2}{R^2} \left(1 - \frac{r^2}{R_n^2} \right), \quad (4)$$

$$\sigma_{T_n} = \frac{\alpha TE}{4(1-\mu)} \times \frac{R^2 - R_n^2}{k^2} \left(3 \frac{r^2}{R_n^2} - 1 \right), \quad (5)$$

Card 2/10

Residual Stresses During Surface....

S/129/61/000/004/010/012

E073/E535

$$\sigma_{oc_n} = \frac{\alpha TE}{2(1-\mu)} \cdot \frac{R^2 - R_n^2}{R^2} \times \left(2 \frac{r^2}{R_n^2} - 1 \right), \quad (6)$$

In these equations σ_r , σ_T and σ_{oc} are, respectively, the radial, tangential and axial stresses, α - coefficient of linear expansion, μ - Poisson coefficient, E - Young's modulus, T - temperature difference between the internal and external layers of the cylinder, R - radius, as shown in Fig.1. In deriving these formulae the following assumptions were made:

1. Cooling is only from the external surface of the specimen;
2. the temperature distribution in the heated layer is symmetrical relative to the axis of the cylinder and is constant along its length;
3. the magnitudes of the axial deformations, which are sufficiently distant from the end faces, are the same and equal the relative axis deformation of a specimen of unit length;
4. the values of μ , α and E in the range of elastic deformations are independent of the temperature.

Card 3/10

Residual Stresses During Surface....

S/129/61/000/004/010/012
E073/E535

Analysis of the above formulae indicates that, during surface hardening of cylindrical components, the stresses in the surface will always consist of axial compression stresses and tangential residual thermal stresses, the magnitude of which depends on the thickness of the heated layer and its temperature. Fig.2 gives the dependence of the maximum tangential thermal residual stresses at the surface of the specimen as a function of F_3/F and αT . G. F. Golovin and M. M. Zamyatnin (Ref.3) have determined experimentally the total residual tangential stresses in 65 mm diameter specimens of Steel 45 after induction hardening to a depth of 6 mm and low temperature tempering; their results are reproduced in Fig.4. The results of approximate analytical and experimental determination of the stresses during surface hardening to a depth of 7 mm are reproduced in Fig.5. In another paper Golovin determined the total residual tangential stresses in a specimen ($R = 32.5$ mm) hardened to a depth of 5.5 mm. He determined the values R_n and R_c at 29 and 24.5 mm, respectively, from the hardness distribution curve along the cross-section. Comparison of the calculated results with experimental data are given in Fig.6. It is concluded that the

Card 4/10


Residual Stresses During Surface.... S/129/61/000/004/010/012
E073/E535

total stresses occurring during induction surface hardening can be determined with adequate accuracy by means of approximate analytical formulae. There are 6 figures and 8 references: all Soviet.

ASSOCIATION: Moskovskiy institut neftekhimicheskoy i gazovoy promyshlennosti (Moscow Institute of the Petrochemical and Gas Industries)

Fig.1. Legend:

Sketch of the temperature distribution along the cross section of a steel cylindrical specimen heated for surface hardening (t_y - temperature at which the metal loses its elastic properties).
 $t_{H04} - t_{initial}$, $R_{H04} - R_{initial}$



Card 5/10


Residual Stresses During Surface....

S/129/61/000/004/010/012
E073/E535

Fig.2 Legend:

Dependence of the maximum tangential residual thermal stresses (σ_T , kg/mm²) at the specimen surface on F_3/F and αT .

$F_3 = \pi (R^2 - R_n^2)$ - area of the layer heated for hardening,
 $F = \pi \cdot R^2$ - total area of the specimen cross section.



Card 6/10


Residual Stresses During Surface...

S/129/61/000/004/010/012

EO73/E535

Fig.3. Legend:

Dependence of the tangential structural stresses (σ_T , kg/mm²) at the specimen surface on the ratio F_3/F and T/M for $\ell_0 = 10^{-3}$.
 M - temperature of the beginning of martensite transformation,
 ℓ_0 - deformation associated with the martensite transformation.



Card 7/10

30

Residual Stresses During Surface....


S/129/61/000/004/010/012
E073/E535

Fig.4. Legend:

Tangential stresses in the case of surface hardening to a depth of 6 mm. σ_T , kg/mm² vs. radius, mm.

- Curve 1 - thermal stresses,
- " 2 - structural stresses,
- " 3 - total stresses (calculated)
- " 4 - curve determined experimentally by G. F. Golovin.

Card 8/10



Residual Stresses During Surface....

S/129/61/000/004/010/012

E073/E535


Fig.5. Legend:

Stresses in the case of surface hardening to a depth of 7 mm.

σ_T , kg/mm² vs. radius, mm.

Curve 1 - calculated

" 2 - experimental



Card 9/10

Residual Stresses During Surface....

S/129/61/000/004/010/012

E073/E535


Fig.6. Legend:

Tangential stresses in the case of surface hardening to a depth of 5.5 mm.

σ_T , kg/mm² vs. radius, mm.

- Curve 1 - thermal stresses,
- " 2 - structural stresses,
- " 3 - total stresses (calculated)
- " 4 - total stresses (experimentally determined).

Card 10/10



KAN, A.A.

New data on root aphids. Uzb. biol. zhur. 6 no.2:64-69 '62.
(MIRA 15:4)

1. Institut zoologii i parazitologii AN UzSSR.
(UZBEKISTAN--PLANT LICE)

KAN, A.G.

Mud solution and its therapeutic value in the treatment of gynecological diseases. Akush. gin., Moskva no.5:83-84 Sept-Oct 1952. (GLML 23:2)

1, Professor.

KAN, A.G.

Thermal stresses and some problems in designing equipment using
two-layer metal. Izv. vys. ucheb. zav.; neft' i gaz 4 no.9:
83-89 '61. (MIRA 14:12)

1. Moskovskiy institut neftekhimicheskoy i gazovoy promyshlennosti
imeni akademika I.M. Gubkina.

(Thermal stresses)

(Petroleum refineries--Equipment and supplies)